

This article was downloaded by: [Academy of Sciences of China]

On: 28 July 2009

Access details: Access Details: [subscription number 907210226]

Publisher Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Journal of Asian Natural Products Research

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t713454007>

New geranyloxycoumarins from *Toddalia asiatica*

Fei Wang^{ab}; Yao Xu^b; Ji-Kai Liu^a

^a State Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, China ^b BioBioPha Co., Ltd, Kunming, China

Online Publication Date: 01 August 2009

To cite this Article Wang, Fei, Xu, Yao and Liu, Ji-Kai(2009)'New geranyloxycoumarins from *Toddalia asiatica*',*Journal of Asian Natural Products Research*,11:8,752 — 756

To link to this Article: DOI: 10.1080/10286020903048975

URL: <http://dx.doi.org/10.1080/10286020903048975>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

New geranyloxycoumarins from *Toddalia asiatica*

Fei Wang^{ab*}, Yao Xu^b and Ji-Kai Liu^{a*}

^aState Key Laboratory of Phytochemistry and Plant Resources in West China, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming 650204, China; ^bBioBioPha Co., Ltd, 132 Lanhei Road, Kunming 650204, China

(Received 1 April 2009; final version received 18 May 2009)

A phytochemical study on the twigs of *Toddalia asiatica* led to the isolation of two new geranyloxycoumarins, 7-geranyloxy-5-methoxycoumarin (**1**) and 8-geranyloxy-5,7-dimethoxycoumarin (**2**). Their structures were elucidated on the basis of extensive spectroscopic analysis.

Keywords: *Toddalia asiatica*; Rutaceae; geranyloxycoumarin

1. Introduction

Toddalia asiatica (L.) Lam. (Rutaceae), a woody liana growing in subtropical and tropical areas of China, has been widely used in traditional Chinese herbal medicine to treat cold, stomachache, intention neuralgia, rheumatism, and injuries from falls [1]. All parts of the plant are claimed to have medicinal value, but the roots, in particular, are believed to be more potent. Decoctions or infusions of the root bark have good efficacy in the treatment of chronic lumbago and sciatica [1]. Previously, several research groups have reported some bioactive components from this medicinal plant, including coumarins [2–7], alkaloids [6,8–10], coumarin–quinolone dimer [11], and coumarin–naphthoquinone dimer [12].

As part of our effort to assemble a natural compound library possessing thousands of structures coming from plants and micro-organisms, further chemical investigation on this plant led to the isolation of two new coumarin–monoterpene ethers,

7-geranyloxy-5-methoxycoumarin (**1**) and 8-geranyloxy-5,7-dimethoxycoumarin (**2**), together with four known coumarins, artanin (**3**) [4], norbraylin (**4**) [2], 5,7,8-trimethoxycoumarin (**5**) [2], and toddalosin (**6**) [5] (Figure 1). Herein, details of the isolation and structural elucidation of compounds **1** and **2** are described.

2. Results and discussion

Compound **1** was obtained as an amorphous powder. Its molecular formula was determined to be C₂₀H₂₄O₄ on the basis of positive ESI-MS at *m/z* 351 [M+Na]⁺, in combination with the ¹³C NMR (DEPT) spectrum. The UV spectrum showed the absorption maxima at 248 (3.90), 257 (3.89), and 329 (4.28) nm, typical of a coumarin chromophore. The ¹H NMR spectrum (Table 1) exhibited characteristic AB system signals of the lactone ring of a coumarin at δ_H 6.15 and 7.96 (each 1H, d, *J* = 9.6 Hz) assigned to H-3 and H-4, respectively. The ¹³C NMR spectrum (Table 1) exhibited 20 carbon signals,

*Corresponding authors. Email: wangfei@mail.kib.ac.cn; jkliu@mail.kib.ac.cn




Figure 1. Structures of compounds **1**–**6**.

including nine carbon signals due to the coumarin skeleton at δ_C 162.9 (s), 161.6 (s), 156.9 (s), 156.7 (s), 138.8 (d), 110.8 (d), 103.9 (s), 95.3 (d), and 93.5 (d), 10 typical geranyloxy carbon resonances at δ_C 142.5 (s), 132.0 (s), 123.6 (d), 118.2 (d), 65.4 (t), 39.5 (t), 26.2 (t), 25.6 (q), 17.7 (q), and 16.7 (q), and one methoxy carbon at δ_C 55.9 (q), indicative of a 5,7-dioxygenated coumarin derivative. The following important HMBC correlations (Table 1) were observed: from the protons at δ_H 7.96 (1H, d, $J = 9.6$ Hz, H-4) and 3.87 (3H, s, OMe) to the carbon at δ_C 156.9 (s, C-5), and from the protons at δ_H 4.57 (2H, d, $J = 6.6$ Hz, H-1') to the carbon at δ_C 162.9 (s, C-7), which confirmed that the methoxy and geranyloxy groups were unambiguously attached at C-5 and C-7, respectively. Therefore, the structure of **1** was elucidated as 7-geranyloxy-5-methoxycoumarin.

Compound **2**, an amorphous powder, possessed a molecular formula of $C_{21}H_{26}O_5$ based on the positive ESI-MS at m/z 381 $[M+Na]^+$ and supported by the ^{13}C NMR (DEPT) spectrum. The UV spectrum also showed the absorption maxima of a coumarin chromophore at 264 (4.00) and 322 (4.07) nm. The NMR spectra (Table 1) were very similar to those of **1**, and there were only two

obvious differences as follows: (1) an additional methoxy signal was observed and (2) one aromatic proton within the coumarin skeleton was absent, suggesting that this compound was a trioxxygenated coumarin substituted by one geranyloxy and two methoxy groups. The following significant HMBC correlations (Table 1): from the protons at δ_H 7.97 (1H, d, $J = 9.5$ Hz, H-4) and 3.91 (3H, s, 5-OMe) to the carbon at δ_C 152.2 (s, C-5), from the protons at δ_H 3.91 (3H, s, 5-OMe) and 3.95 (3H, s, 7-OMe) to the carbon at δ_C 91.2 (d, C-6), and from the protons at δ_H 4.60 (2H, d, $J = 7.0$ Hz, H-1') to the carbon at δ_C 128.7 (s, C-8) were detected, which indicated that two methoxy groups and one geranyloxy must be linked at C-5, C-7, and C-8, respectively. Accordingly, the structure of **2** was characterized as 8-geranyloxy-5,7-dimethoxycoumarin.

3. Experimental

3.1 General experimental procedures

UV spectra were obtained on a Shimadzu UV-2401PC spectrophotometer. IR spectra were performed on a Bruker Tensor 27 FT-IR spectrometer with KBr pellets. 1D and 2D NMR spectra were respectively recorded on Bruker AV-400 and DRX-500

Table 1. ¹H and ¹³C NMR spectroscopic data for compounds **1** and **2** in CDCl₃.

No.	1			2		
	δ _H	δ _C	HMBC	δ _H	δ _C	HMBC
2	—	161.6 (s)		—	160.9 (s)	
3	6.15 (1H, d, 9.6)	110.8 (d)	2, 4a	6.15 (1H, d, 9.5)	111.2 (d)	2, 4a
4	7.96 (1H, d, 9.6)	138.8 (d)	5, 8 ^a , 8a	7.97 (1H, d, 9.5)	138.7 (d)	5, 8 ^a , 8a
4a	—	103.9 (s)		—	103.9 (s)	
5	—	156.9 (s)		—	152.2 (s)	
6	6.30 (1H, d, 1.3)	95.3 (d)	4a, 8	6.32 (1H, s)	91.2 (d)	4a, 8
7	—	162.9 (s)		—	156.5 (s)	
8	6.41 (1H, d, 1.3)	93.5 (d)	4a, 6	—	128.7 (s)	
8a	—	156.7 (s)		—	149.1 (s)	
1'	4.57 (2H, d, 6.6)	65.4 (t)	7, 3'	4.60 (2H, d, 7.0)	69.9 (t)	8, 3'
2'	5.46 (1H, br t, 6.6)	118.2 (d)	4', 5'	5.58 (1H, br t, 7.0)	119.7 (d)	4', 5'
3'	—	142.5 (s)		—	142.5 (s)	
4'	2.09 (2H, m)	39.5 (t)	2', 5', 7'	2.02 (2H, m)	39.6 (t)	2', 5', 7'
5'	1.76 (3H, s)	16.7 (q)	2', 4'	1.67 (3H, s)	16.4 (q)	2', 4'
6'	2.13 (2H, m)	26.2 (t)	3', 8'	2.05 (2H, m)	26.4 (t)	3', 8'
7'	5.08 (1H, br t, 5.9)	123.6 (d)	4', 9', 10'	5.06 (1H, br t, 6.0)	123.9 (d)	
8'	—	132.0 (s)		—	131.6 (s)	
9'	1.67 (3H, br s)	25.6 (q)	7', 10'	1.66 (3H, br s)	25.7 (q)	7', 10'
10'	1.60 (3H, br s)	17.7 (q)	7', 9'	1.58 (3H, br s)	17.7 (q)	7', 9'
OCH ₃	3.87 (3H, s, 5-OMe)	55.9 (q)	5, 6 ^a	3.91 (3H, s, 5-OMe)	56.0 (q)	5, 6 ^a
				3.95 (3H, s, 7-OMe)	56.4 (q)	6 ^a , 7

^aWeak but significant four-bond HMBC correlations.

instruments with the deuterated solvent (CDCl_3 : δ_{H} 7.26, δ_{C} 77.0) as an internal standard. EI-MS and ESI-MS (including HR-ESI-MS) were measured on Finnigan-MAT 90 and API QSTAR Pulsar i mass spectrometers, respectively. Silica gel (200–300 mesh; Qingdao Marine Chemical, Inc., Qingdao, China) and Sephadex LH-20 (Amersham Biosciences, Uppsala, Sweden) were used for column chromatography (CC). Fractions were monitored by TLC (Qingdao Marine Chemical, Inc.) in combination with Agilent 1200 HPLC (Eclipse XDB-C18 column, 5 μm , 4.6×150 mm, 50–100% MeOH in H_2O over 8 min followed by 100% MeOH to 15 min, 1 ml/min, 25°C).

3.2 Plant material

The twigs of *T. asiatica* were collected from Yunnan Province, China and identified by Prof. Dr Hua Peng. The voucher specimen (No. BBP2009004TA) has been deposited in the Herbarium of Kunming Institute of Botany, Chinese Academy of Sciences.

3.3 Extraction and isolation

The air-dried and powdered twigs (5.0 kg) were extracted with 95% ethanol at room temperature and filtered. The filtrate was concentrated to give a crude extract (~400 g), which was subjected to silica gel CC with a stepwise elution of petroleum ether (PE)–acetone to yield fractions A–I. Fraction B (~5.0 g) eluted by PE–acetone (96:4) was further separated by silica gel CC using PE–acetone (100:1, 50:1, 10:1, 1:1) to yield subfractions A–D, respectively. Subfraction A (~300 mg) was repeatedly isolated and purified by silica gel CC (PE–acetone = 150:1; PE– CHCl_3 = 5:1) and Sephadex LH-20 (CHCl_3 –MeOH = 1:1), and recrystallized to afford compound **1** (91 mg). Similarly, compound **2** (296 mg) was

obtained from subfraction B (~750 mg) by silica gel CC (PE–acetone = 50:1; PE– CHCl_3 = 3:1) and Sephadex LH-20 (CHCl_3 –MeOH = 1:1).

3.3.1 7-Geranyloxy-5-methoxycoumarin (**1**)

An amorphous powder; HPLC: t_{R} = 10.24 min; UV λ_{max} (CHCl_3) (log ϵ): 248 (3.90), 257 (3.89), 329 (4.28) nm; IR (KBr) ν_{max} : 3075, 3016, 1725, 1630, 1612, 1501, 1452, 1364, 1226, 1196, 1164, 1123 cm^{-1} ; NMR spectral data: see Table 1; EI-MS m/z : 192 [M –geranyl] $^{+}$ (100), 164 (39), 149 (11), 136 (19); ESI-MS (pos.) m/z : 351 [M +Na] $^{+}$; HR-ESI-MS (pos.) m/z : 351.1569 (calcd for $\text{C}_{20}\text{H}_{24}\text{O}_4\text{Na}$, 351.1572).

3.3.2 8-Geranyloxy-5,7-dimethoxycoumarin (**2**)

An amorphous powder; HPLC: t_{R} = 9.25 min; UV λ_{max} (CHCl_3) (log ϵ): 264 (4.00), 322 (4.07) nm; IR (KBr) ν_{max} : 3084, 1730, 1605, 1505, 1438, 1345, 1261, 1221, 1188, 1149, 1120 cm^{-1} ; NMR spectral data: see Table 1; EI-MS m/z : 222 [M –geranyl] $^{+}$ (100), 207 (34), 193 (26), 179 (9), 136 (6); ESI-MS (pos.) m/z : 381 [M +Na] $^{+}$; HR-ESI-MS (pos.) m/z : 381.1676 (calcd for $\text{C}_{21}\text{H}_{26}\text{O}_5\text{Na}$, 381.1677).

Acknowledgements

This work was financially supported by the National Basic Research Program of China (973 Program) 2009CB522300 and the R&D funds from BioBioPha Co., Ltd.

References

- [1] Editorial Committee of Flora Reipublicae Popularis Sinicae, *Flora Reipublicae Popularis Sinicae* (Academic Press, Beijing, 1997), Vol. 43 (2), p. 96.
- [2] M.N. Deshmukh, V.H. Deshpande, and A.V.R. Rao, *Phytochemistry* **15**, 1419 (1976).

- [3] P.N. Sharma, A. Shoeb, R.S. Kapil, and S.P. Popli, *Phytochemistry* **20**, 335 (1981).
- [4] G. Szabó, H. Greger, and O. Hofer, *Phytochemistry* **24**, 537 (1985).
- [5] H. Ishii, J.I. Kobayashi, K.I. Yamaguchi, and T. Ishikawa, *Chem. Pharm. Bull.* **41**, 1655 (1993).
- [6] I.L. Tsai, M.F. Wun, C.M. Teng, T. Ishikawa, and I.S. Chen, *Phytochemistry* **48**, 1377 (1998).
- [7] H.A. Oketch-Rabah, J.W. Mwangi, J. Lisgarten, and E.K. Mberu, *Fitoterapia* **71**, 636 (2000).
- [8] I.S. Chen, I.L. Tsai, S.J. Wu, W.S. Sheen, T. Ishikawa, and H. Ishii, *Phytochemistry* **34**, 1449 (1993).
- [9] I.L. Tsai, S.C. Fang, T. Ishikawa, C.T. Chang, and I.S. Chen, *Phytochemistry* **44**, 1383 (1997).
- [10] S.C. Jain, M.K. Pandey, R.K. Upadhyay, R. Kumar, G. Hundal, and M.S. Hundal, *Phytochemistry* **67**, 1005 (2006).
- [11] H. Ishii, J.I. Kobayashi, and T. Ishikawa, *Tetrahedron Lett.* **32**, 6907 (1991).
- [12] H. Ishii, J.I. Kobayashi, H. Seki, and T. Ishikawa, *Chem. Pharm. Bull.* **40**, 1358 (1992).